



# **NUECES BBASC STUDY #3**

## **NUECES WATERSHED PRE- AND POST-DEVELOPMENT NUTRIENT BUDGETS**

**NUECES ESTUARY ADVISORY COUNCIL**

**JUNE 22, 2015**

**PAULA JO LEMONDS, PE, PG**

# DISCUSSION

Background

Status

Results

Schedule

# BACKGROUND

- Nueces BBASC work plan
  - Tier 2b Recommendation
- Nueces BBEST
  - BBEST Recommendations Report
    - Sec. 5.2 Nutrient Considerations
- Nueces BBASC
  - BBASC Recommendations Report
    - Sec. 4.3.2 Nutrient Considerations

# GOALS

- Develop nutrient budgets based on quantitative understanding of natural supply of all nutrient forms and anthropogenic changes in these supplies over time for Nueces Bay watershed
- Determine annual loads for both the pre-development and present condition

Source: Nueces BBASC work plan



# SCOPE OF WORK

- Task 1 – Compile Data
  - Compile Water Quality and Hydrologic Data
- Task 2 – Perform Data Evaluation and Modeling Analyses
  - Perform Data Evaluation
    - Available data, sampling stations, and subwatersheds
    - Identify appropriate stations
  - Refine Linear Regression Analyses
  - Estimate Reservoir Influence Sink/Sources
  - Water Quality Correlations to Anthropogenic Changes
  - Pre- and Post-Development Loadings







# ANALYSES

- Land Use
- Mission-Aransas Watershed
- Wastewater Treatment Plant Effects
- Effects of CCR Construction
- Annual Load Calculation for Dry, Average and Wet Years; Pre- and Post-CCR Construction









# LAND USE ANALYSIS

- Using NLCD to analyze land use changes over time



# LAND USE




- About 4% increase in urban area over 1992-2011
- Cultivated acres relatively steady




# MISSION-ARANSAS WATERSHED ANALYSIS

- Very little development in watershed
- Hypothesized that M-A watershed would provide an additional evaluation as to whether changes in land use could be influencing water quality.



# MISSION-ARANSAS WATERSHED ANALYSIS

- Upper Mission River had higher nutrient concentrations than upstream stations in less developed portions of Nueces
  - However,  $\text{NO}_x$  higher in Nueces compared to upper Mission River
  - Where comparisons possible, observed nutrient loadings in upper Aransas were even higher than observed in Mission River and upstream stations in the Nueces.
- One explanation for difference in nutrient loadings is land use in Aransas River subwatershed.
  - Land use is predominantly hay/pasture and cultivated crops








# PLEASANTON EXAMPLE


- Station 12981, Atascosa River at Pleasanton
- $\text{NH}_4$ ,  $\text{NO}_x$ , TP,  $\text{PO}_4$  and perhaps Chl-a higher at 12981 than 12980
- NRA BSR (2013a) states that much of upper Atascosa River is intermittent or ephemeral
  - If flows were not augmented by effluent from the Pleasanton WWTP outfall, river might be classified as intermittent, except for lowermost segment.






# UPSTREAM TO DOWNSTREAM

- 12981 – Pleasanton
- 12980 – Downstream



Regression analysis, Station 12981

Nueces Tributaries, Texas (1970-2014)



Regression analysis, Station 12980


Nueces Tributaries, Texas (1970-2014)

# DETERMINING THE EFFECTS OF CCR CONSTRUCTION


- Assumed data pre-1986 represented pre-construction and data post-1986 represented post-construction period
  - N species decline between pre- and post- reservoir construction
    - TKN decline may have begun prior to construction
  - TP and PO<sub>4</sub>: Some increases in TP and PO<sub>4</sub> in the Frio River and San Miguel Creek post-construction that do not appear downstream of the reservoir at Three Rivers
  - Flow: Low flow conditions below CCR are greater than before CCR due to flow management

# EFFECTS OF CCR - TKN


- TKN data from three TCEQ locations
  - Frio River at Tilden, Station 13023
  - San Miguel Creek nr Tilden, Station 12983
  - Three Rivers, Station 12979



Nutrient Distributions, Station 13023



Nutrient Distributions, Station 12983




Nutrient Distributions, Station 12979


● Observed Concentration  
● Estimated Concentration

# EFFECTS OF CCR - TOTAL PHOSPHORUS


- Nutrient data from two TCEQ locations at
  - Frio River, Station ID 13023 (Figure 7-1) and
  - San Miguel Creek, Station ID 12983 (Figure 7-2)
  - Three Rivers, Station 12979



Nutrient Distributions, Station 13023



Nutrient Distributions, Station 12983



Nutrient Distributions, Station 12979

● Observed Concentration  
● Estimated Concentration

# ANNUAL LOAD CALCULATIONS

- $\text{Load} = \text{Concentration} \times \text{Flow}$
- Concentration time-series
- Dry, average, wet years of USGS streamflow




Representative flow years based on  
Nueces at Mathis (USGS 08211000, TCEQ 12965)

	Dry	Average	Wet
Pre-1986	1984	1974	1971
Post-1986	2008	1993	2002

- Indication of nutrient load delivered to Estuary
- Also looked at dry, avg, wet years based on Calallen precip statistics

# CONCENTRATION TIME-SERIES

## Nutrient Distributions, Station 12979 Nueces Tributaries, Texas (1959-2014)







- Observed Concentration
- Estimated Concentration




# LOADS

- TP, TN, TKN, NO<sub>x</sub>



# EPA ECOREGION REFERENCE CONDITIONS



Parameter	EPA 25th Percentile Reference Conditions			
(mg/L)	Ecoregion IV, Subcoregion 30 <sup>1</sup>	Ecoregion IV, Subcoregion 31 <sup>2</sup>	Ecoregion IX, Subcoregion 33 <sup>3</sup>	Ecoregion X, Subcoregion 34 <sup>4</sup>
Total Phosphorus	0.008	0.028	0.1	0.126
Total Kjedaahl Nitrogen (TKN)	0.18	0.27	0.543	0.74
Total Nitrogen	0.27*, 0.55 <sup>†</sup>	0.49*	0.681*, 0.935 <sup>†</sup>	0.88*, 0.86 <sup>†</sup>
Nitrite + Nitrate - N	0.09	0.22	0.138	0.14
Chlorophyll a**	0.002	0.002	0.000733	0.0021
Turbidity (FTU)	0.73	3.83	10.9	12.27
* Calculated				
† Reported				
** Chlorophyll a measured by Spectrophotometric method with acid correction				

# CONCLUSIONS

- Land use slowly changing. Urban areas increasing in size. Areas of cultivated crops seem to remain steady (1970 to present).
- Some locations are affected by WWTP discharge.
- Effects of CCR Construction
  - N species decline between pre- and post- reservoir construction
    - TKN decline may have begun prior to the construction
  - TP and  $\text{PO}_4$ : Appear to be some increases in TP and  $\text{PO}_4$  in the Frio River and San Miguel Creek post-construction that do not appear downstream of reservoir at Three Rivers
  - Flow: Low flow conditions below CCR are greater than before CCR due to flow management
- Difficult to determine statistically significant relationships between flow and nutrients
  - Dataset is representative of a range of conditions.
  - Correlations could improve if they were done for specific temporal periods, rising or declining parts of hydrograph, or seasonally.

# RECOMMENDATIONS

- Evaluate predictive scenarios of loadings upstream and downstream of CCR with CCR/LCC System operating under different operational schemes
- Evaluate effects of other development scenarios, including future build-out land use conditions, on nutrient loadings
- Quantify effects of seasonality on nutrient loadings.
- Further quantify effects of reservoirs in Nueces Watershed.
  - Fate and transport, nutrient processing uptake, losses, releases, dissolved oxygen
- Identify and quantify nonpoint source component of nutrient loadings in Nueces Watershed
  - One way to accomplish quantification of nonpoint source loadings is with watershed-scale nutrient loading model that takes into account point sources, as well as nonpoint sources.

# TASK 3 – MEETINGS AND REPORT







- Task 3 – Meetings and Report
  - NEAC Kickoff Meeting June 16, 2014
  - NEAC Meeting Update October 20, 2014
  - NEAC Meeting Update February 23, 2015
  - NEAC Meeting Update June 22, 2015
  - Draft Report: June 30, 2015
  - Final Report, Contract Deadline: August 31, 2015







- Station 12999- Figure 4-14
- Station 12965-Appendix, Page 52
- Station 12979-Figure 4-15
- Station 12980-Figure 6-6
- Station 13024-Appendix, Page 61



Regression analysis, Station 12979

Nueces Tributaries, Texas (1970-2014)

USGS_8190000 - Upstream Gage (unmanaged)				USGS_8211000 - Downstream Gage (heavily managed)				Calallen Precip gage (suburb of Corpus)			
average:	63,413		65,217	average:	300,975		174,663	average:	35		30
				geo mean	212,152	geo mean	106,381				
1970-1985 sorted		1986-2014 sorted		1970-1985 sorted		1986-2014 sorted		1970-1985 sorted		1986-2014 sorted	
1980	16,470	2011	7,332	1984	47,248	2008	37,988	1982	22.77	2011	12.66
1978	27,440	2014	8,406	1983	54,263	2009	39,873	1977	26.55	1989	16.65
1983	28,543	2013	14,342	1982	105,443	2006	42,512	1984	26.89	1996	19.01
1979	32,156	2009	17,004	1978	112,263	2013	43,355	1974	27.58	1988	19.33
1984	33,268	2012	17,169	1972	150,368	2011	43,506	1975	29.69	2012	19.51
1982	36,012	1989	19,500	1970	180,665	2014	48,476	1978	31.55	2009	20.84
1974	50,317	2006	22,038	1979	183,261	2012	50,937	1980	32.61	2008	23.15
1975	50,855	1993	23,700	1975	188,417	2010	54,589	1970	33.92	2013	25
1970	56,310	2008	26,594	1974	196,536	1988	57,763	1972	34.69	2005	25.08
1972	64,369	2010	31,279	1985	236,812	2000	58,231	1979	37.14	2000	25.9
1985	68,317	1995	33,684	1977	266,578	1996	58,523	1983	37.87	2014	27.08
1977	78,523	1988	36,329	1980	283,516	1989	60,802	1985	39.65	2006	27.54
1976	84,524	1994	40,524	1976	467,987	1986	64,335	1971	40.71	1999	27.87
1973	116,091	2005	56,935	1973	525,632	1995	77,821	1976	43.55	1990	27.97
1981	133,250	2003	57,389	1981	532,307	1991	92,005	1973	45.46	1986	29.51
1971	138,161	1999	58,364	1971	1,284,308	1994	93,991	1981	45.76	2001	32.14
		2002	58,460			1999	100,190			1987	32.33
		1986	63,850			1993	101,551			2003	33.03
		2000	84,151			2005	113,366			1995	33.11
		1990	85,830			1997	125,382			1994	33.73
		1996	86,082			2001	130,139			1993	34.95
		1991	97,526			1998	168,851			1998	36
		2001	111,889			1990	177,480			2004	38.56
		1998	121,648			2003	256,412			2002	39.34
		1992	128,554			1987	381,824			1991	41.58
		1997	131,474			2004	452,732			1992	41.82
		1987	146,386			1992	466,299			2007	42.59
		2007	149,315			2007	539,718			1997	43.12
		2004	155,525			2002	1,126,578			2010	43.55